

The environmental costs and benefits of high-yield farming

Andrew Balmford^{1*}, Tatsuya Amano^{1,2}, Harriet Bartlett¹, Dave Chadwick³, Adrian Collins⁴, David Edwards⁵, Rob Field⁶, Philip Garnsworthy⁷, Rhys Green¹, Pete Smith⁸, Helen Waters¹, Andrew Whitmore⁹, Donald M. Broom¹⁰, Julian Chara¹¹, Tom Finch^{1,6}, Emma Garnett¹, Alfred Gathorne-Hardy^{12,13,14}, Juan Hernandez-Medrano¹⁵, Mario Herrero¹⁶, Fangyuan Hua¹, Agnieszka Latawiec^{17,18}, Tom Misselbrook⁴, Ben Phalan^{1,19}, Benno I. Simmons¹, Taro Takahashi^{4,20}, James Vause²¹, Erasmus zu Ermgassen¹ and Rowan Eisner¹

How we manage farming and food systems to meet rising demand is pivotal to the future of biodiversity. Extensive field data suggest that impacts on wild populations would be greatly reduced through boosting yields on existing farmland so as to spare remaining natural habitats. High-yield farming raises other concerns because expressed per unit area it can generate high levels of externalities such as greenhouse gas emissions and nutrient losses. However, such metrics underestimate the overall impacts of lower-yield systems. Here we develop a framework that instead compares externality and land costs per unit production. We apply this framework to diverse data sets that describe the externalities of four major farm sectors and reveal that, rather than involving trade-offs, the externality and land costs of alternative production systems can covary positively: per unit production, land-efficient systems often produce lower externalities. For greenhouse gas emissions, these associations become more strongly positive once forgone sequestration is included. Our conclusions are limited: remarkably few studies report externalities alongside yields; many important externalities and farming systems are inadequately measured; and realizing the environmental benefits of high-yield systems typically requires additional measures to limit farmland expansion. Nevertheless, our results suggest that trade-offs among key cost metrics are not as ubiquitous as sometimes perceived.

Agriculture already covers around 40% of Earth's ice- and desert-free land and is responsible for around two-thirds of freshwater withdrawals¹. Its immense scale means that it is already the largest source of threat to other species², so how we cope with very marked increases in demand for farm products^{3,4} will have profound consequences for the future of global biodiversity^{2,5}. On the demand side, cutting food waste and excessive consumption of animal products is essential^{1,5-8}. In terms of supply, farming at high yields (production per unit area) has considerable potential to restrict humanity's impacts on biodiversity. Detailed field data from five continents and almost 1,800 species from birds to daisies⁹⁻¹⁴ reveal that so many depend on native vegetation that for most the impacts of agriculture on their populations would be best limited by farming at high yields (production per unit area) alongside sparing large tracts of intact habitat. Provided it can be coupled with setting aside (or restoring) natural habitats¹⁵, lowering the land cost of agriculture thus appears central to addressing the extinction crisis².

However, a key counterargument against this land-sparing approach is that there are many other environmental costs of agriculture besides the biodiversity displaced by the land it requires, such as greenhouse gas (GHG) and ammonia emissions, soil erosion, eutrophication, dispersal of harmful pesticides and freshwater depletion^{5,7,16-18}. Measured per unit area of farmland, the production of such externalities is sometimes greater in high- than lower-yield farming systems^{17,18}, potentially weakening the case for land sparing. However, while expressing externalities per unit area can help identify local-scale impacts¹⁹, it systematically underestimates the overall impact of lower-yield systems that occupy more land for the same level of production²⁰. To be robust, assessments of externalities also need to include the off-site effects of management practices, such as crop production for supplementary feeding of livestock, or off-farm grazing for manure inputs to organic systems²⁰⁻²².

¹Conservation Science Group, Department of Zoology, University of Cambridge, Cambridge, UK. ²Centre for the Study of Existential Risk, University of Cambridge, Cambridge, UK. ³Environment Centre Wales, Bangor, UK. ⁴Rothamsted Research, Okehampton, UK. ⁵Department of Animal and Plant Sciences, University of Sheffield, Sheffield, UK. ⁶RSPB Centre for Conservation Science, The Royal Society for the Protection of Birds, Sandy, UK. ⁷School of Biosciences, University of Nottingham, Loughborough, UK. ⁸Institute of Biological and Environmental Sciences, University of Aberdeen, Aberdeen, UK. ⁹Rothamsted Research, Harpenden, UK. ¹⁰Department of Veterinary Medicine, University of Cambridge, Cambridge, UK. ¹¹CIPAV, Centre for Research on Sustainable Agricultural Production Systems, Cali, Colombia. ¹²School of Geosciences, University of Edinburgh, Edinburgh, UK. ¹³Global Academy of Agriculture and Food Security, University of Edinburgh, Edinburgh, UK. ¹⁴Oxford India Centre for Sustainable Development, Somerville College, Oxford, UK. ¹⁵Faculty of Veterinary Medicine and Zootechny, National Autonomous University of Mexico, Mexico City, Mexico. ¹⁶Commonwealth Scientific and Industrial Research Organisation, St Lucia, Queensland, Australia. ¹⁷Department of Geography and Environment, Pontifical Catholic University of Rio de Janeiro (PUC-Rio), Rio de Janeiro, Brazil. ¹⁸Institute of Agricultural Engineering and Informatics, Faculty of Production and Power Engineering, University of Agriculture in Kraków, Kraków, Poland. ¹⁹Universidade Federal da Bahia, Rua Barão de Jeremoabo, Salvador, Brazil. ²⁰British Veterinary School, University of Bristol, Bristol, UK. ²¹UN Environment World Conservation Monitoring Centre, Cambridge, UK. *e-mail: a.balmford@zoo.cam.ac.uk




Fig. 1 | Framework for exploring how different environmental costs compare across alternative production systems. **a**, A hypothetical plot of externality cost versus land cost of different, potentially interchangeable production systems (blue circles) in a given farming sector. In this example, the data suggest a trade-off between externality and land costs across different systems. **b**, This example reveals a more complex pattern, with additional systems (in green and red circles) that are low or high in both costs.

A novel framework for comparing system-wide costs

Here, we argue that comparisons of the overall impacts of contrasting agricultural systems should focus on the sum of externality generated per unit of production¹⁰ (paralleling measures of emissions intensity in climate change analyses). This approach has, for the most part, been adopted only for a relatively narrow set of agricultural products^{8,23} and farming systems (for example, organic versus conventional, glasshouse versus open-field^{20,24}). Here we develop a more general framework, and apply it to a diversity of data on some major farm sectors, farming systems and environmental externalities. Existing data are limited but nevertheless enable us to explore the utility of this new approach, test for broad patterns and make an informed commentary on their significance for understanding the trade-offs and co-benefits of high- versus lower-yield systems.

Our framework involves plotting the environmental costs of producing a given quantity of a commodity against one another, across alternative production systems (as in Fig. 1). We focus on examining variation in some better-known externality costs in relation to land cost (that is, $1/\text{yield}$), because of the latter's fundamental importance as a proxy for impacts on biodiversity. However, the approach could be used to explore associations among any other costs for which data are available. Comparisons must be made across production systems that could, in principle, be substituted for one another, so they must be measured or modelled identically and in the same place or, if not, potential confounding effects of different methods, climate and soils must be removed statistically. If the idea that high-yield systems impose disproportionate externalities is true, we would expect plots of externality per unit production against land cost to show negative associations (Fig. 1a, blue symbols). However observed patterns may be more complex, and could reveal promising systems associated with low land cost and low externalities, or unpromising systems with high land and externality costs (Fig. 1b, green and red symbols, respectively).

Our team of sector and externality specialists collated data for applying this framework to five major externalities (GHG emissions, water use, nitrogen (N), phosphorus (P) and soil losses) in four major sectors (Asian paddy rice, European wheat, Latin American beef and European dairy; Methods). We used both literature searches and consultation with experts to find paired yield and externality measurements for contrasting production systems in each sector. To be included, data had to be near-complete for a given externality—for example, most major elements of GHG emissions or N losses had to be included, and if systems involved inputs (such as feeds or fertilizers) generated off-site we required

data on the externality and land costs of their production. To limit confounding effects, we narrowed our geographic scope within each sector (Supplementary Table 1), so that differences across systems could reasonably be attributed to farm practices rather than gross bioclimatic variation. Where co-products were generated, we apportioned overall costs among products using economic allocation, but also investigated alternative allocation rules.

Findings for four sectors

Our first key result is that useable data are surprisingly scarce. Few studies measured paired externality and yield information, many reported externalities in substantially incomplete or irreconcilably divergent ways, and we could find no suitable data at all on some widely adopted practices. Nevertheless, we were able to obtain sufficient data to consider how externalities vary with land costs for 9 out of 20 possible sector–externality combinations (Supplementary Table 1). The type of data available differed across these combinations (which we view as a useful test of the flexibility of our framework). For one combination, the most extensive data we could find was from a long-term experiment at a single location. However because we were interested in generalities, where possible we used information from multiple studies—either field experiments or life cycle assessments (LCAs) conducted across several sites—and used generalized linear mixed models (GLMMs) to correct for confounding method and site effects (Methods). Last, for two sectors, we used process-based models parameterized for a fixed set of conditions representative of the region.

The data that we were able to obtain do not suggest that environmental costs are generally larger for farming systems with low land costs (that is, high-yield systems; Fig. 2). If anything, positive associations—in which high-yield, land-efficient systems also have lower costs in other dimensions—appear more common. For Chinese paddy rice, we found sufficient multi-site experimental data to explore how two focal externalities vary with land cost across contrasting systems (Methods). GHG costs (Fig. 2a) showed negative associations with land cost across monoculture and rotational systems (assessed separately). Our GLMMs revealed that for both system types, greater application of organic N lowered land cost but increased emissions (probably because of feedstock effects on the methanogenic community²⁵; Supplementary Table 2); by contrast, there was little or no GHG penalty from boosting yield using inorganic N (arrows, Fig. 2a). A large volume of data on rice and water use showed weakly positive covariation in costs (Fig. 2b). GLMMs indicated that increasing application of inorganic N boosted




Fig. 2 | Externality costs of alternative production systems against land cost for five externalities in four agricultural sectors. All costs are expressed per tonne of production (so land cost, for instance, is in ha-years per tonne; that is, the inverse of yield). **a–j**, Costs differentiated by sector (shown by icons). **a, b**, Asian paddy rice. **c, d**, European wheat. **e, f**, Latin American beef. **g–j**, European dairy. Different externalities are indicated by background shading (grey, GHG emissions; blue, water use; pink, N emissions; purple, P emissions; buff, soil loss), and different sectors (Asian paddy rice, European wheat, Latin American beef and European dairy) are shown by icons. Points on plots derived from multi-site experiments (**a–c**) and LCAs (**e**) show values for systems adjusted for site and study effects via GLMMs of land cost and externality cost (for 95% confidence intervals, see Supplementary Fig. 1), while arrows show management practices with statistically significant effects (whose 95% confidence intervals do not overlap zero in the GLMMs; Methods). In **d** (wheat and N emissions), progressively darker circles depict increasing nitrate application rate (0, 48, 96, 144, 192, 240 and 288 kg N per ha-year). In **f** (beef and GHG emissions, estimated by RUMINANT), different colours show different system types. In **g–j** (dairy and four externalities), circles and squares show results for conventional and organic systems, respectively (detailed in Supplementary Table 4). Spearman’s rank correlation coefficients (P values) are **a**, rice–rice: -0.51 (0.002); rice–cereal, -0.36 (0.06); **b**, 0.19 (0.26); **c**, -0.34 (0.14); **d**, -0.21 (0.66); **e**, 0.95 (0.001); **f**, 0.83 (<0.001); **g**, 0.90 (0.08); **h**, 0.70 (0.23); **i**, 1.00 (0.02); and **j**, 1.00 (0.02). Note that these correlation coefficients do not necessarily reflect nonlinear relationships (for example, **d**) accurately. Credit: Icons for Asian paddy rice and European wheat: Freepik (www.flaticon.com).

yield²⁶, and less irrigation lowered water use while incurring only a modest yield penalty²⁷ (Supplementary Table 2). Sensitivity tests of the rice analyses had little impact on these patterns (Methods; Supplementary Fig. 2).

We found two useable data sets on European wheat, both from the UK (Methods). Our GLMMs of data from a three-site experiment varying the N fertilization regime revealed a complex relationship between GHG and land costs (Fig. 2c; Supplementary Table 2), driven by divergent responses²⁸ to adding ammonium nitrate (which lowers land costs but increases embodied GHG emissions) and adding urea (which lowers land costs without increasing GHG emissions per unit production, but at the cost of increased ammonia volatilization). A single-site experiment varying inorganic N treatments showed a nonlinear relationship between land cost and N losses (Fig. 2d), with increasing N application lowering both costs until an apparent threshold, beyond which land cost decreased further but at the cost of greater N leaching (see also ref.¹).

In livestock systems, all data we could find showed positive covariation between land costs and externalities. For Latin American beef, we located coupled yield estimates only for GHG emissions, but here two different types of data (Methods) revealed a common pattern. Using GLMMs again to control for potentially confounding study and site effects, we found that across multiple LCAs, pasture systems with greater land demands also generated greater emissions (Fig. 2e), with both land and GHG costs reduced by pasture improvements (using N fertilization or legumes). This pattern across contrasting pasture systems was confirmed by running RUMINANT²⁹ (Fig. 2f), a process-based model that also identified relatively low land and GHG costs for a series of silvopasture and feedlot-finishing systems (for which comparable LCA data were unavailable).

For European dairy, process-based modelling of three conventional and two organic systems, parameterized for the UK, enabled us to estimate four different externalities alongside yield (Methods). This showed that conventional systems—especially those using less grazing and more concentrates—had substantially lower land and also GHG costs (Fig. 2g), in part because concentrates reduce CH₄ emissions from fibre digestion³⁰. Systems with greater use of concentrates (which have less rumen-degradable protein than grass³¹) also showed lower losses of N, P and soil per unit production (Fig. 2h–j). These broad patterns persisted when we used protein production rather than economic value to allocate costs to co-products (Methods; Supplementary Fig. 2).

Incorporating land use

As a final analysis, we examined the additional externalities resulting from the different land requirements of contrasting systems. To generate the same quantity of agricultural product, low-yield systems require more land, allowing less to be retained or restored as natural habitat. This is in turn likely to increase GHG emissions and soil loss, and alter hydrology—although we could find only enough data to explore the first of these effects. For each sector, we supplemented our direct GHG figures for each system with estimates of GHG consequences of their land use following Intergovernmental Panel on Climate Change (IPCC) methods³² to calculate the sequestration potential of a hectare not used for farming and instead allowed to revert to climax vegetation (Methods). Results (Fig. 3) showed that these GHG opportunity costs of agriculture were typically greater than the emissions from farming activities themselves and, when added to them, in every sector generated strongly positive cross-system associations between overall GHG cost and land cost. These patterns were maintained in sensitivity tests where we halved recovery rates or assumed half of the area potentially freed from farming was retained under agriculture (Methods; Supplementary Fig. 3). These findings thus confirm recent suggestions^{33,34} that high-yield farming has the potential, provided land not needed for production

is largely used for carbon sequestration, to make a substantial contribution to mitigating climate change.

Conclusions, caveats and knowledge gaps

This study was conceived as an exploration of whether high-yield systems—central to the idea of sparing land for nature in the face of enormous human demand for farm products—typically impose greater negative externalities than alternative approaches. Our results support three conclusions. First, useful data are worryingly limited. We considered only four relatively well-studied sectors and a narrow set of externalities—not including important impacts such as soil health or the effects of pesticide exposure on human health²⁰. Even then, we found studies reporting yield-linked estimates of externalities scarce, with many widely adopted or promising practices within these sectors undocumented. We were not able to examine complex agricultural systems (such as mixed farming or agroforestry) that might have relatively low externalities. Relevant data on many significant developing-world farm sectors (such as cassava or dryland cereal production in Africa) also appear very limited. Given that a multi-dimensional understanding of the environmental effects of alternative production systems is integral to delivering sustainable intensification, more field measurements linking yield with a broader suite of externalities across a much wider range of practices and sectors are urgently needed.

Second, the available data on the sector–externality combinations that we considered do not suggest that negative associations between land cost and other environmental costs of farming are typical (in contrast to Fig. 1a). Many low-yield systems impose high costs in other ways too and, although certain yield-improving practices have undesirable impacts (for example, organic fertilization of paddy rice increasing CH₄ emissions; see also ref.¹), other practices appear capable of reducing several costs simultaneously (see also refs^{1,8,24,35,36}). High (but not excessive) application of inorganic N, for example, can lower land take of Chinese rice production without incurring GHG or water-use penalties. Similarly, in Brazilian beef production, adopting better pasture management, semi-intensive silvopasture and feedlot-finishing can all boost yields alongside lowering GHG emissions. It is worth noting that although most systems we examined are relatively high-yielding, other recent work suggests that positive associations (rather than trade-offs) among environmental and land costs may, if anything, be more likely in lower-yielding systems¹.

Third, pursuing promising high-yield systems is clearly not the same as encouraging business-as-usual industrial agriculture. Some high-yield practices we did not examine, such as the heavy use of pesticides in much tropical fruit cultivation³⁷, are likely to increase externality costs per unit production. Of the high-yield practices we did investigate some, such as applying fossil-fuel-derived ammonium nitrate to UK wheat, impose disproportionately high environmental costs. Others that seem favourable in terms of our focal externalities incur other costs, such as high NH₃ emissions from using urea on wheat²⁸, and management regimes that reduce costs in one geographic setting may not do so in others¹. Much work characterizing existing systems and designing new ones is thus needed. We suggest that our framework can serve as a device for identifying existing yield-enhancing systems that also lower other environmental costs—and perhaps more importantly, for benchmarking the environmental performance of promising new technologies and practices.

We close by stressing that for high-yield systems to generate any environmental benefits they must be coupled with efforts to reduce rebound effects. Several plausible mechanisms for limiting these by explicitly linking yield growth to improved environmental performance have been identified—including strict land-use zoning; strategic deployment of yield-enhancing loans, expertise or infrastructure; conditional access to markets; and restructured




Fig. 3 | Overall GHG cost against land cost of alternative systems in each sector, including the GHG opportunity costs of land under farming. The y-axis values are the sum of GHG emissions from farming activities (plotted in Fig. 2a,c,e,g) and the forgone sequestration potential of land maintained under farming and thus unable to revert to natural vegetation (Methods). All costs are expressed per tonne of production. **a–d**, GHG costs differentiated by sector (shown by icons). **a**, Asian paddy rice. **b**, European wheat. **c**, Latin American beef. **d**, European dairy. The notation is as in Fig. 2. Spearman's rank correlation coefficients (P values) are **a**, rice-rice: 0.40 (0.017); rice-cereal: 0.80 (<0.001); **b**, 0.99 (<0.001); **c**, 0.98 (<0.001); and **d**, 0.80 (0.13). Credit: Icons for Asian paddy rice and European wheat: Freepik (www.flaticon.com).

rural subsidies¹⁵. Without such linkages, systems that perform well per unit production may nevertheless cause net environmental harm through higher profits or lower prices stimulating land conversion^{38–40}, and damage human health by encouraging overconsumption of cheap, calorie-rich but nutrient-deficient foods^{41,42}. If promising high-yield strategies are to help solve rather than exacerbate society's challenges, yield increases instead need to be combined with far-reaching demand-side interventions^{1,6,41} and directly linked with effective measures to constrain agricultural expansion¹⁵.

Methods

Focal sectors and externalities. We focused on 4 globally significant farm sectors (Asian paddy rice, European wheat, Latin American beef and European dairy, accounting for 90%, 33%, 23% and 53% of global output of these products⁴³) and 5 major externalities (GHG emissions, water use, N, P and soil losses). We chose these sector–externality combinations because preliminary work suggested that they were characterized quantitatively relatively often, using diverse approaches (single-site experiments, multi-site experiments, LCAs and process-based models), enabling us to explore the generality of our framework. We then searched the literature and consulted experts to obtain paired yield and externality estimates of alternative production systems in each sector, narrowing our geographic scope so that differences in system performance could be reasonably attributed to management practices (rather than gross variation in bioclimate or soils). Our analyses have rarely been attempted previously and have complex data requirements, so we could not adopt standard procedures developed for systematic reviews on topics where many studies have attempted to answer the same research question.

This process generated data on ≥ 5 contrasting production systems for 9 out of 20 possible sector–externality combinations (Supplementary Table 1): Chinese rice–GHG emissions (from multi-site experiments); Chinese rice–water use

(multi-site experiments); UK wheat–GHG emissions (a multi-site experiment); UK wheat–N emissions (a single-site experiment); Brazilian beef–GHG emissions (both LCA data and process-based models); and UK dairy–GHG emissions, and N, P and soil losses (process-based models). Water use in the wheat and most of the beef systems examined was limited and so not explored further. We could not find sufficient paired yield–externality estimates for the nine remaining sector–externality combinations.

The land and externality costs of each system were then expressed as total area used per unit production (that is, $1/\text{yield}$) and total amount of externality generated per unit production. All estimates included the area used and externalities generated in producing externally derived inputs (such as feed or fertilizers). For analytical tractability, as in other recent studies^{1,24}, we treat impacts occurring at different times and places as being additive. Occasional gaps in estimates for a system were filled using standard values from IPCC or other sources, or information from study authors or comparable systems (details below). Where experiments or LCAs were conducted at multiple sites, we built GLMMs in the package lme4⁴⁴ in R version 3.3.1⁴⁵ to identify effects of specific management practices on land and externality cost estimates adjusted for potentially confounding biophysical and methodological effects. To illustrate the effects of statistically significant management variables (those whose 95% confidence intervals did not overlap zero; shown in bold in Supplementary Table 2), we estimated land and externality costs at the observed minimum and maximum values (for continuous management variables) or with the reference category and the category that showed the maximum effect size (for categorical variables), while keeping other variables constant; we then linked these points as arrows on our externality cost/land cost plots (Fig. 2 and Supplementary Figs. 1 and 2, with arrows displaced horizontally and/or vertically for increased visibility). Where systems generated significant co-products (wheat and rapeseed from rotational rice, beef from dairy), we allocated land and externality costs to the focal product in proportion to its relative contribution to the gross monetary value of production per unit area of farmland (from focal and co-product combined)⁴⁶.

Rice and GHG emissions. Systematic searching of Scopus for experimental studies reporting both yields and emissions of Chinese paddy rice systems identified 17 recently published studies^{47–63} containing 140 paired yield–emissions estimates for different systems (after within-year replicates of a system were averaged). To limit confounding effects, we analysed separately the data from monoculture systems from southern provinces (2 rice crops per year; 5 studies, 60 estimates) and rotational systems from more northerly provinces (1 rice and 1 wheat or rape crop per year; 12 studies, 80 estimates). The studies documented the effects of variation in tillage (yes/no), application rates of inorganic and organic N, and (only for rotational systems) irrigation regime (continuous flooding versus episodic midseason drainage). There were insufficient data to examine effects of seedling density, crop variety, organic practices, biochar application, use of groundcover to lower emissions, N fertilizer type, or K or P fertilization.

Land cost estimates were expressed in ha-years per tonne of rice grain (that is, the inverse of annual production per hectare farmed). GHG costs were expressed in tonnes of CO₂ equivalents (CO₂e) per tonne of rice grain, and included CH₄ and N₂O emissions for growing and fallow seasons (with the latter where necessary based on mean values from refs^{47–49,64}), and embodied emissions from N fertilizer production (Yara emissions database; F. Brendrup, personal communication). We were unable to include emissions from producing manure or K or P fertilizer, or from farm machinery. For rotational systems, we adjusted the land and GHG costs of rice production downwards by multiplying them by the proportional contribution of rice to the gross monetary value of production per unit area of farmland from rice and co-product combined (using mean post-2000 prices from ref.⁴³).

We next built GLMMs predicting variation in our estimates of land cost and GHG cost, for the monoculture and rotational data sets in turn. Management practices assessed as predictors were tillage regime (binary), application rates of organic N and of inorganic N, and irrigation regime (binary; only rotational systems). Study site was included as a random effect. For all systems, we adjusted for biophysical and methodological differences across sites using the first two components from a principal component analysis of site scores for 14 variables: annual precipitation, precipitation during the driest and wettest quarters, annual mean temperature, mean temperatures during the warmest and coldest quarters, maximum temperature during the warmest month, mean monthly solar radiation, latitude, longitude, soil organic carbon content, plot size, replicates per estimate, and start year (with all climate data taken from refs^{65,66}). Principal components 1 and 2 together explained 82.3% and 76.2% of the variance in these variables for monoculture and rotational systems, respectively. Soil pH and (soil pH)² were also assessed as additional predictors. For the monoculture models, tolerance values were all >0.4 (indicating an absence of multicollinearity) except for the pH terms (both <0.1), which we therefore removed. For the rotational models, all tolerance values indicated an absence of multicollinearity, but (soil pH)² was removed because AICc (corrected Akaike Information Criterion) values indicated that model fit was no better than using soil pH alone. Final models (Supplementary Table 2) were then used to plot site-adjusted land and GHG costs (as points) and statistically significant management effects (as arrows) in Fig. 2a. We also tested the effect of allocating land and GHG costs in rotational systems based on the relative energy content of rice and co-products⁶⁷ (rather than a relative contribution to gross monetary value; Supplementary Fig. 2).

We adopted similar although simpler approaches for the next two sector–externality combinations, which again used data from multi-site experiments.

Rice and water use. A systematic search on Scopus yielded 15 recent studies^{57,58,64,68–79} meeting our criteria containing 123 paired estimates describing the effects of variation in inorganic N application rate and irrigation regime on land and water costs of Chinese paddy rice. We analysed monoculture and rotational systems together but considered water use solely for periods of rice production. Land cost was expressed in ha-years per tonne of rice grain, and water cost in cubic metres per tonne of rice grain (excluding rainfall). We adjusted these estimates for site effects in GLMMs of variation in land and water costs using as predictors the application rate of inorganic N, and irrigation regime (a six-level factor: continuous flooding, continuous flooding with drainage, alternate wetting and drying, controlled irrigation, mulches or plastic films, and long periods of dry soil), while accounting for the effect of study site as a random effect. Tolerance values were all >0.7. Final models (Supplementary Table 2) were then used to plot site-adjusted land and water costs (points) and significant management effects (arrows) in Fig. 2b. Almost all sources reported data on only one rice season per year, but one study⁶⁸ included separate estimates for early- and late-season rice, so we checked the robustness of our findings by re-running the analysis without the early-season data from this study (Supplementary Fig. 2).

Wheat and GHG emissions. The Agricultural Greenhouse Gas Inventory Research Platform^{80–83} provided 96 paired measures of variation in yield and N₂O emissions in response to experimental changes in N fertilizer application rate and type. We expanded the emissions profile to include embodied emissions from N fertilizer production (from the Yara emissions database; F. Brendrup, personal communication). We derived land costs in ha-years per tonne of wheat (at 85% dry matter) and GHG costs in tonnes of CO₂e per tonne of wheat. Experiments

were run in three regions, so to adjust for site effects we built GLMMs of variation in land and GHG costs fitting study region as a random effect and using the application rates of ammonium nitrate, urea and dicyandiamide (a nitrification inhibitor) as predictors. Tolerance values were all >0.7. Adjusted land and GHG cost estimates from the final models (Supplementary Table 2) are plotted in Fig. 2c, with arrows showing statistically significant management practices.

Wheat and N losses. We assessed this sector–externality combination using data from Rothamsted's long-term Broadbalk wheat experiment, which investigates the effects of inorganic N application rates on yields of winter wheat. During the 1990s changes in field drainage enabled the measurement (alongside yield) of plot-specific leaching losses of nitrate⁸⁴. Mean land and N costs—expressed in ha-years per tonne of wheat (at 85% dry matter) and kilograms of N leached per tonne of wheat, respectively—were averaged across 8 seasons (thus smoothing-out rainfall effects), for each of 7 levels of N application (from 0 to 288 kg N (as ammonium nitrate) per ha-year; details in Fig. 2 legend). Results are plotted in Fig. 2d.

Beef and GHG emissions. Two types of data were available for this sector–externality combination, enabling us to compare findings across assessment techniques. First we examined all published LCAs of Brazilian beef production^{85–92}. Supplementing this with a bioclimatically comparable data set from tropical Mexico (R. Olea-Perez, personal communication) yielded 33 paired yield–emissions estimates for contrasting production systems. These varied in whether they used improved pasture, supplementary feeding or improved breeds (which if unreported we inferred from age at first calving, and mortality and conception rates). There were insufficient LCA data to examine the effects of feedlots, silvopasture or rotational grazing. Land costs were calculated in ha-years per tonne of carcass weight (CW), incorporating land used to grow feed, and assuming a dressing percentage of 50%⁹³. GHG costs were derived in tonnes of CO₂e per tonne of CW, including enteric CH₄ emissions, CH₄ and N₂O emissions from manure, N₂O emissions from managed pasture, emissions from supplementary feed production (where necessary using values from ref.⁸⁶), and embodied GHG emissions from N, P and K fertilizer production. There were too few data to include CO₂ emissions from lime application or farm machinery. Milk production was not a significant co-product. To control for site effects, we built GLMMs of variation in land and GHG costs using site as a random effect and use of improved pasture, supplementary feeding and improved breeds (each a binary factor) as predictors. Tolerance values were all >0.8. Adjusted land and GHG cost estimates from the final models (Supplementary Table 2) are plotted in Fig. 2e, with the arrow describing a statistically significant management practice.

For comparison we derived an equivalent GHG cost versus land cost plot (Fig. 2f) using a process-based model of beef production. RUMINANT²⁹ is an IPCC tier 3 digestion and metabolism model that uses stoichiometric equations to estimate production of meat, manure N and enteric methane for any given pasture quality, supplementary feed quantity and type, cattle breed and region. We used plausible combinations of these settings (Supplementary Table 3) and corresponding values of feed and forage protein, digestibility and carbohydrate content (judged representative of the Brazilian beef sector by M.H.) to derive yield and emissions estimates for 86 contrasting pasture systems. To extend beyond the scope of the LCA analyses, we also modelled 50 silvopasture systems by boosting feed quality to simulate access to *Leucaena*, and 8 feedlot-finishing systems by incorporating an 83–120 day feedlot phase when animals received high-quality mixed ration. For each system, we included the whole herd, after determining the ratio of fattening/breeding animals using the DYNMOD demographic projection tool⁹⁴, based on system-specific reproductive performance parameters and animal growth rates (reflecting pasture quality and management; Supplementary Table 3). Breeding animals experienced the same conditions as fattening animals (except that in pasture and silvopasture they received no supplementary feed). Stocking rates were set to sustainable carrying capacity for pasture and silvopasture, and 201 animals ha⁻¹ for feedlots (D.M.B. personal observation). Yields were converted to land cost in ha-years per tonne of CW, including the area of feedlots and land required to grow feed (using feed composition and yield data from refs^{43,85}). RUMINANT emissions estimates were supplemented with estimates of manure CH₄, CO₂ and N₂O emissions from feed production, and N₂O emissions from pasture fertilization (from refs^{32,85}). Carbon sequestration by vegetation could not be included, so we probably overestimate net GHG emissions from silvopasture⁹⁵. All emissions were converted to CO₂e units (using conversion factors from refs^{32,85} and feedlot manure distribution from ref.⁹⁶) and expressed in tonnes of CO₂e per tonne of CW.

Dairy and four externalities. We also used process-based models to investigate how GHG emissions and N, P and soil losses varied with land cost across five dairy systems representative of UK practices (Supplementary Table 4; Fig. 2g–j). We modelled three conventional systems with animals accessing grazing for 270, 180 and 0 days per year, and two organic systems with grazing access for 270 and 200 days per year. Model farms were assigned rainfall and soil characteristics based on frequency distributions of these parameters for real farms of each type, with structural and management data (for example, ratios of livestock categories and ages, N and P excretion rates) based on the models of refs^{31,97,98}. Manure

management was based on representative variations of the 'manure management continuum'⁹⁹ (Supplementary Table 4). Physical performance data (annual milk yield, concentrate feed input, replacement rate and stocking rate) were obtained from the Agriculture and Horticulture Development Board (AHDB) Dairy database (M. Topliff, personal communication) for conventional systems and from the Department for Environment, Food and Rural Affairs (DEFRA)¹⁰⁰ for organic systems.

Yields were converted to land cost in ha-years per tonne of energy-corrected milk (ECM), including land required to grow feed (from refs^{101,102}, with yield penalties for organic production from ref.¹⁰³). As 57% of global beef production originates from the dairy sector¹⁰⁴, we adjusted land costs downwards by multiplying them by the proportional contribution of milk to the gross monetary value of production per unit area of farmland from milk and beef combined (using prices from the AHDB Dairy database (M. Topliff, personal communication)).

GHG cost estimates for each system comprised CH₄ emissions from enteric fermentation (based on ref.³¹), CH₄ and N₂O emissions from manure management (following refs^{32,105}), emissions from N fertilizer applications to pasture (from refs^{106,107}), and from feed production (from ref.¹⁰⁸). Emissions from farm machinery and buildings were not included. Emissions were then summed and expressed in tonnes of CO₂e per tonne of ECM. Nitrate losses of each system were derived from the National Environment Agricultural Pollution–Nitrate (NEAP-N) model^{109,110}, while P and soil losses were estimated using the Phosphorus and Sediment Yield Characterisation In Catchments (PSYCHIC) model^{98,111}. These last three costs were expressed in kilograms per tonne of ECM and (as with land costs) downscaled by allocating a portion of them to beef co-products, based on milk and beef prices. Finally, to check the effect of this allocation rule, we re-ran each analysis instead allocating costs using the relative protein content of milk and beef (from ref.¹⁰⁴; Supplementary Fig. 2).

GHG opportunity costs of land farmed. Alongside the GHG emissions generated by agricultural activities themselves (analysed above), farming typically carries an additional GHG cost. Wherever the carbon content of farmed land is less than that of the natural habitat that could replace it if agriculture ceased, farming imposes an opportunity cost of sequestration forgone¹¹², whose magnitude increases with the area under production (and hence with the land cost of the system). We quantified this GHG cost using the forgone sequestration method, whereby retaining the current land use is assumed to prevent the sequestration in soils and biomass that would occur if the land was allowed to revert to climax vegetation (see details in Supplementary Table 5).

For each forgone transition, values for annual biomass accrual (≤ 20 years) were taken from Table 4.9 of ref.³², assuming that the climax vegetation for UK wheat and dairy was 'temperate oceanic forest (Europe)'; for Chinese rice it was 'tropical moist deciduous forest (Asia, continental)', and for Brazilian beef it was 'tropical moist deciduous forest (South America)'. The carbon content of all biomass was assumed to be 47% of dry matter (ref.³² Table 4.3).

Changes in soil carbon values were taken from the relevant mean percentage change in soil organic carbon values for each land conversion from a global meta-analysis¹¹³. For UK wheat and Chinese rice, we used values for conversion of cropland to woodland; for UK dairy and Brazilian beef, we used conversion of grassland to woodland for grazing land and conversion of cropland to woodland for land used to grow feed. Initial soil carbon values were taken from Table 2.3 of ref.³². We assumed that the soils for UK wheat were 'cold temperate, moist, high activity soils', for Chinese rice, they were 'tropical, wet, low activity soils', for UK dairy, they were 'cold temperate, moist, high activity soils' for grazing land and for producing imported feed they were 'subtropical humid, LAC soils' (South America), and for Brazilian beef for both grazing and feed production they were 'tropical, moist, low activity soils'. In each case, the relevant percentage change in soil organic carbon was multiplied by the initial soil carbon stock to calculate an absolute change, which, following IPCC guidelines³², we assumed took 20 years.

Total annual forgone sequestration was then estimated by adding this annual change in soil organic carbon and the annual accrual of biomass carbon under reversion to climax vegetation. We assumed (as in ref.³⁴) that each 1 ha reduction in land cost results in 1 ha of recovering habitat. As above, our land cost estimates included land needed to produce externally derived inputs, and (for rotational rice and dairy) were adjusted downwards based on the value of co-products. These GHG opportunity costs were then added to the direct GHG emissions estimates of each system, and the summed values were plotted against land cost (Fig. 3).

As a sensitivity test of our key assumptions, we re-ran these analyses assuming that carbon recovery rates are halved, or that (because of rebound or similar effects^{38–40}) half of the area potentially freed from farming is retained under agriculture. These two changes to our assumptions have numerically identical effects, shown in Supplementary Fig. 3. Note that our recovery-based estimates of the GHG costs that farming imposes through land use are conservative, in that they are roughly 30–50% of those obtained from calculating GHG emissions from natural habitat clearance (annualized, for consistency with the recovery method, over 20 harvests; data not shown).

Code availability. The R codes used for the analyses are available from the corresponding author upon request.

Data availability

The data that support the findings of this study are available from the corresponding author upon request.

Received: 3 April 2018; Accepted: 10 August 2018;
Published online: 14 September 2018

References

- Poore, J. & Nemecek, T. Reducing food's environmental impacts through producers and consumers. *Science* **360**, 987–992 (2018).
- Green, R. E., Cornell, S. J., Scharlemann, J. P. W. & Balmford, A. Farming and the fate of wild nature. *Science* **307**, 550–555 (2005).
- Tilman, D., Balzer, C., Hill, J. & Befort, B. L. Global food demand and the sustainable intensification of agriculture. *Proc. Natl Acad. Sci. USA* **108**, 20260–20264 (2011).
- Hunter, M. C., Smith, R. G., Schipanski, M. E., Atwood, L. W. & Mortensen, D. A. Agriculture in 2050: recalibrating targets for sustainable intensification. *Bioscience* **67**, 386–391 (2017).
- Godfray, H. C. J. et al. Food security: the challenge of feeding 9 billion people. *Science* **327**, 812–818 (2010).
- Bajželj, B. et al. Importance of food-demand management for climate mitigation. *Nat. Clim. Change* **4**, 924–929 (2014).
- Foley, J. A. et al. Solutions for a cultivated planet. *Nature* **478**, 337–342 (2011).
- Ripple, W. J. et al. Ruminants, climate change and climate policy. *Nat. Clim. Change* **4**, 2–5 (2014).
- Phalan, B., Onial, M., Balmford, A. & Green, R. E. Reconciling food production and biodiversity conservation: land sharing and land sparing compared. *Science* **333**, 1289–1291 (2011).
- Balmford, A., Green, R. & Phalan, B. Land for food & land for nature? *Daedalus* **144**, 57–75 (2015).
- Hulme, M. F. et al. Conserving the birds of Uganda's banana-coffee arc: land sparing and land sharing compared. *PLoS ONE* **8**, e54597 (2013).
- Kamp, J. et al. Agricultural development and the conservation of avian biodiversity on the Eurasian steppes: a comparison of land-sparing and land-sharing approaches. *J. Appl. Ecol.* **52**, 1578–1587 (2015).
- Dotta, G., Phalan, B., Silva, T. W., Green, R. & Balmford, A. Assessing strategies to reconcile agriculture and bird conservation in the temperate grasslands of South America: grasslands conservation and agriculture. *Conserv. Biol.* **30**, 618–627 (2016).
- Williams, D. R. et al. Land-use strategies to balance livestock production, biodiversity conservation and carbon storage in Yucatán, Mexico. *Glob. Change Biol.* **23**, 5260–5272 (2017).
- Phalan, B. et al. How can higher-yield farming help to spare nature? *Science* **351**, 450–451 (2016).
- Pretty, J. Agricultural sustainability: concepts, principles and evidence. *Phil. Trans. R. Soc. B* **363**, 447–465 (2008).
- Matson, P. A., Parton, W. J., Power, A. G. & Swift, M. J. Agricultural intensification and ecosystem properties. *Science* **277**, 504–509 (1997).
- Tilman, D., Cassman, K. G., Matson, P. A., Naylor, R. & Polasky, S. Agricultural sustainability and intensive production practices. *Nature* **418**, 671–677 (2002).
- Didham, R. K. et al. Agricultural intensification exacerbates spillover effects on soil biogeochemistry in adjacent forest remnants. *PLoS ONE* **10**, e0116474 (2015).
- Seufert, V. & Ramankutty, N. Many shades of gray – the context-dependent performance of organic agriculture. *Sci. Adv.* **3**, e1602638 (2017).
- Kirchmann, H., Bergström, L., Kätterer, T., Andrén, O. & Andersson, R. in *Organic Crop Production – Ambitions and Limitations* (eds Kirchmann, H. & Bergström, L.) 39–72 (Springer, Dordrecht, 2008).
- Madhusudan, M. D. The global village: linkages between international coffee markets and grazing by livestock in a South Indian wildlife reserve. *Conserv. Biol.* **19**, 411–420 (2005).
- Nijdam, D., Rood, T. & Westhoek, H. The price of protein: review of land use and carbon footprints from life cycle assessments of animal food products and their substitutes. *Food Policy* **37**, 760–770 (2012).
- Clark, M. & Tilman, D. Comparative analysis of environmental impacts of agricultural production systems, agricultural input efficiency, and food choice. *Environ. Res. Lett.* **12**, 64016 (2017).
- Yan, X., Yagi, K., Akiyama, H. & Akimoto, H. Statistical analysis of the major variables controlling methane emission from rice fields. *Glob. Change Biol.* **11**, 1131–1141 (2005).
- Pittelkow, C. M., Adviento-Borbe, M. A., van Kessel, C., Hill, J. E. & Linquist, B. A. Optimizing rice yields while minimizing yield-scaled global warming potential. *Glob. Change Biol.* **20**, 1382–1393 (2014).
- Carrijo, D. R., Lundy, M. E. & Linquist, B. A. Rice yields and water use under alternate wetting and drying irrigation: a meta-analysis. *Field Crop Res.* **203**, 173–180 (2017).

28. Smith, K. A. et al. The effect of N fertilizer forms on nitrous oxide emissions from UK arable land and grassland. *Nutr. Cycl. Agroecosyst.* **93**, 127–149 (2012).
29. Herrero, M. et al. Biomass use, production, feed efficiencies, and greenhouse gas emissions from global livestock systems. *Proc. Natl Acad. Sci. USA* **110**, 20888–20893 (2013).
30. Beauchemin, K., McAllister, T. A. & McGinn, S. M. Dietary mitigation of enteric methane from cattle. *CAB Rev. Perspect. Agric. Vet. Sci. Nutr. Nat. Resour.* **4**, 1–18 (2009).
31. Wilkinson, J. M. & Garnsworthy, P. C. Dietary options to reduce the environmental impact of milk production. *J. Agric. Sci.* **155**, 334–347 (2017).
32. IPCC 2006 IPCC Guidelines for National Greenhouse Gas Inventories (eds Eggleston, H. S. et al.) (IGES, 2006).
33. Gilroy, J. J. et al. Optimizing carbon storage and biodiversity protection in tropical agricultural landscapes. *Glob. Change Biol.* **20**, 2162–2172 (2014).
34. Lamb, A. et al. The potential for land sparing to offset greenhouse gas emissions from agriculture. *Nat. Clim. Change* **6**, 488–492 (2016).
35. Cui, Z. et al. Pursuing sustainable productivity with millions of smallholder farmers. *Nature* **555**, 363–366 (2018).
36. Notarnicola, B. et al. The role of life cycle assessment in supporting sustainable agri-food systems: a review of the challenges. *J. Clean. Prod.* **140**, 399–409 (2017).
37. Bravo, V. et al. Monitoring pesticide use and associated health hazards in Central America. *J. Int. J. Occup. Environ. Heal.* **173**, 1077–3525 (2011).
38. Lambin, E. F. & Meyfroidt, P. Global land use change, economic globalization, and the looming land scarcity. *Proc. Natl Acad. Sci. USA* **108**, 3465–3472 (2011).
39. Ewers, R. M., Scharlemann, J. P. W., Balmford, A. & Green, R. E. Do increases in agricultural yield spare land for nature? *Glob. Change Biol.* **15**, 1716–1726 (2009).
40. Byerlee, D., Stevenson, J. & Villoria, N. Does intensification slow crop land expansion or encourage deforestation? *Glob. Food Sec.* **3**, 92–98 (2014).
41. Tilman, D. & Clark, M. Global diets link environmental sustainability and human health. *Nature* **515**, 518–522 (2014).
42. Yang, Q. et al. Added sugar intake and cardiovascular diseases mortality among US adults. *JAMA Intern. Med.* **174**, 516 (2014).
43. FAOSTAT: Food and Agriculture Data (Food and Agriculture Organization of the United Nations, 2017); <http://fao.org/faostat>
44. Bates, D., Mächler, M., Bolker, B. & Walker, S. Fitting linear mixed-effects models using lme4. *J. Stat. Softw.* **67**, 1–48 (2015).
45. R: A Language and Environment for Statistical Computing (R Foundation for Statistical Computing, 2016); <https://www.r-project.org>
46. Guinée, J. B., Heijungs, R. & Huppes, G. Economic allocation: examples and derived decision tree. *Int. J. Life Cycle Assess.* **9**, 23–33 (2004).
47. Shang, Q. et al. Net annual global warming potential and greenhouse gas intensity in Chinese double rice-cropping systems: a 3-year field measurement in long-term fertilizer experiments. *Glob. Change Biol.* **17**, 2196–2210 (2011).
48. Liu, Y. et al. Net global warming potential and greenhouse gas intensity from the double rice system with integrated soil-crop system management: a three-year field study. *Atmos. Environ.* **116**, 92–101 (2015).
49. Chen, Z., Chen, F., Zhang, H. & Liu, S. Effects of nitrogen application rates on net annual global warming potential and greenhouse gas intensity in double-rice cropping systems of the Southern China. *Environ. Sci. Pollut. Res. Int.* **23**, 24781–24795 (2016).
50. Xue, J. F. et al. Assessment of carbon sustainability under different tillage systems in a double rice cropping system in Southern China. *Int. J. Life Cycle Assess.* **19**, 1581–1592 (2014).
51. Shen, J. et al. Contrasting effects of straw and straw-derived biochar amendments on greenhouse gas emissions within double rice cropping systems. *Agric. Ecosyst. Environ.* **188**, 264–274 (2014).
52. Ma, Y. C. et al. Net global warming potential and greenhouse gas intensity of annual rice-wheat rotations with integrated soil-crop system management. *Agric. Ecosyst. Environ.* **164**, 209–219 (2013).
53. Zhang, X., Xu, X., Liu, Y., Wang, J. & Xiong, Z. Global warming potential and greenhouse gas intensity in rice agriculture driven by high yields and nitrogen use efficiency. *Biogeosciences* **13**, 2701–2714 (2016).
54. Yang, B. et al. Mitigating net global warming potential and greenhouse gas intensities by substituting chemical nitrogen fertilizers with organic fertilization strategies in rice-wheat annual rotation systems in China: a 3-year field experiment. *Ecol. Eng.* **81**, 289–297 (2015).
55. Zhang, Z. S., Guo, L. J., Liu, T. Q., Li, C. F. & Cao, C. G. Effects of tillage practices and straw returning methods on greenhouse gas emissions and net ecosystem economic budget in rice-wheat cropping systems in central China. *Atmos. Environ.* **122**, 636–644 (2015).
56. Xiong, Z. et al. Differences in net global warming potential and greenhouse gas intensity between major rice-based cropping systems in China. *Sci. Rep.* **5**, 17774 (2015).
57. Xu, Y. et al. Improved water management to reduce greenhouse gas emissions in no-till rapeseed-rice rotations in Central China. *Agric. Ecosyst. Environ.* **221**, 87–98 (2016).
58. Xu, Y. et al. Effects of water-saving irrigation practices and drought resistant rice variety on greenhouse gas emissions from a no-till paddy in the central lowlands of China. *Sci. Total Environ.* **505**, 1043–1052 (2015).
59. Yao, Z. et al. Nitrous oxide and methane fluxes from a rice-wheat crop rotation under wheat residue incorporation and no-tillage practices. *Atmos. Environ.* **79**, 641–649 (2013).
60. Xia, L., Wang, S. & Yan, X. Effects of long-term straw incorporation on the net global warming potential and the net economic benefit in a rice-wheat cropping system in China. *Agric. Ecosyst. Environ.* **197**, 118–127 (2014).
61. Zhang, A. et al. Change in net global warming potential of a rice-wheat cropping system with biochar soil amendment in a rice paddy from China. *Agric. Ecosyst. Environ.* **173**, 37–45 (2013).
62. Zou, J., Huang, Y., Zong, L., Zheng, X. & Wang, Y. Carbon dioxide, methane, and nitrous oxide emissions from a rice-wheat rotation as affected by crop residue. *Adv. Atmos. Sci.* **21**, 691–698 (2004).
63. Zhou, M. et al. Nitrous oxide and methane emissions from a subtropical rice-rapeseed rotation system in China: a 3-year field case study. *Agric. Ecosyst. Environ.* **212**, 297–309 (2015).
64. Yao, Z. et al. Improving rice production sustainability by reducing water demand and greenhouse gas emissions with biodegradable films. *Sci. Rep.* **7**, 39855 (2017).
65. Hijmans, R. J., Cameron, S. E., Parra, J. L., Jones, P. G. & Jarvis, A. *WorldClim – Global Climate Data: WorldClim Version 2* (2017); <http://www.worldclim.org/version2>
66. Hijmans, R. J., Cameron, S. E., Parra, J. L., Jones, P. G. & Jarvis, A. *WorldClim – Global Climate Data: Bioclimatic Variables* (2017); <http://www.worldclim.org/bioclim>
67. Heuzé, V., Tran, G. & Hassoun, P. *Feedipedia: Rough Rice (Paddy Rice)* (Feedipedia, a programme by INRA, CIRAD, AFZ and FAO, 2015); <https://www.feedipedia.org/node/226>
68. Liang, K. et al. Grain yield, water productivity and CH₄ emission of irrigated rice in response to water management in south China. *Agric. Water Manag.* **163**, 319–331 (2016).
69. Kreye, C. et al. Fluxes of methane and nitrous oxide in water-saving rice production in north China. *Nutr. Cycl. Agroecosyst.* **77**, 293–304 (2007).
70. Lu, W., Cheng, W., Zhang, Z., Xin, X. & Wang, X. Differences in rice water consumption and yield under four irrigation schedules in central Jilin Province, China. *Paddy Water Environ.* **14**, 473–480 (2016).
71. Jin, X. et al. Water consumption and water-saving characteristics of a ground cover rice production system. *J. Hydrol.* **540**, 220–231 (2016).
72. Sun, H. et al. CH₄ emission in response to water-saving and drought-resistance rice (WDR) and common rice varieties under different irrigation managements. *Water Air Soil Pollut.* **227**, 47 (2016).
73. Wang, X. et al. The positive impacts of irrigation schedules on rice yield and water consumption: synergies in Jilin Province, Northeast China. *Int. J. Agric. Sustain.* **14**, 1–12 (2016).
74. Xiong, Y., Peng, S., Luo, Y., Xu, J. & Yang, S. A paddy eco-ditch and wetland system to reduce non-point source pollution from rice-based production system while maintaining water use efficiency. *Environ. Sci. Pollut. Res.* **22**, 4406–4417 (2015).
75. Shao, G.-C. et al. Effects of controlled irrigation and drainage on growth, grain yield and water use in paddy rice. *Eur. J. Agron.* **53**, 1–9 (2014).
76. Liu, L. et al. Combination of site-specific nitrogen management and alternate wetting and drying irrigation increases grain yield and nitrogen and water use efficiency in super rice. *Field Crop Res.* **154**, 226–235 (2013).
77. Chen, Y., Zhang, G., Xu, Y. J. & Huang, Z. Influence of irrigation water discharge frequency on soil salt removal and rice yield in a semi-arid and saline-sodic area. *Water* **5**, 578–592 (2013).
78. Ye, Y. et al. Alternate wetting and drying irrigation and controlled-release nitrogen fertilizer in late-season rice. Effects on dry matter accumulation, yield, water and nitrogen use. *Field Crop Res.* **144**, 212–224 (2013).
79. Peng, S. et al. Integrated irrigation and drainage practices to enhance water productivity and reduce pollution in a rice production system. *Irrig. Drain.* **61**, 285–293 (2012).
80. Bell, M. J. et al. Nitrous oxide emissions from fertilised UK arable soils: fluxes, emission factors and mitigation. *Agric. Ecosyst Environ.* **212**, 134–147 (2015).
81. Bell, M. J. et al. *Agricultural Greenhouse Gas Inventory Research Platform – InveN2Ory: Fertiliser Experimental Site in East Lothian, 2011 Version: 1* [data set] (Freshwater Biological Association, 2017); <https://doi.org/10.17865/ghgno606>
82. Cardenas, L. M., Webster, C. & Donovan, N. *Agricultural Greenhouse Gas Inventory Research Platform – InveN2Ory: Fertiliser Experimental Site in*

- Bedfordshire, 2011 Version: 1 [data set] (Freshwater Biological Association, 2017); <https://doi.org/10.17865/ghgno613>
83. Williams, J. R. et al. *Agricultural Greenhouse Gas Inventory Research Inventory Research Platform - InveN2Ory: Fertiliser Experimental Site in Herefordshire, 2011 Version: 1* [data set] (Freshwater Biological Association, 2017); <https://doi.org/10.17865/ghgno675>
 84. Goulding, K. W. T., Poulton, P. R., Webster, C. P. & Howe, M. T. Nitrate leaching from the Broadbalk Wheat Experiment, Rothamsted, UK, as influenced by fertilizer and manure inputs and the weather. *Soil Use Manag.* **16**, 244–250 (2000).
 85. Cardoso, A. S. et al. Impact of the intensification of beef production in Brazil on greenhouse gas emissions and land use. *Agric. Syst.* **143**, 86–96 (2016).
 86. de Figueiredo, E. B. et al. Greenhouse gas balance and carbon footprint of beef cattle in three contrasting pasture-management systems in Brazil. *J. Clean. Prod.* **142**, 420–431 (2017).
 87. Dick, M., Abreu Da Silva, M. & Dewes, H. Life cycle assessment of beef cattle production in two typical grassland systems of southern Brazil. *J. Clean. Prod.* **96**, 426–434 (2015).
 88. Florindo, T. J., de Medeiros Florindo, G. I. B., Talamini, E., da Costa, J. S. & Ruviaro, C. F. Carbon footprint and life cycle costing of beef cattle in the Brazilian midwest. *J. Clean. Prod.* **147**, 119–129 (2017).
 89. Mazzetto, A. M., Feigl, B. J., Schils, R. L. M., Cerri, C. E. P. & Cerri, C. C. Improved pasture and herd management to reduce greenhouse gas emissions from a Brazilian beef production system. *Livest. Sci.* **175**, 101–112 (2015).
 90. Pashaei Kamali, F. et al. Environmental and economic performance of beef farming systems with different feeding strategies in southern Brazil. *Agric. Syst.* **146**, 70–79 (2016).
 91. Ruviaro, C. F., De Léis, C. M., Lampert, V. D. N., Barcellos, J. O. J. & Dewes, H. Carbon footprint in different beef production systems on a southern Brazilian farm: a case study. *J. Clean. Prod.* **96**, 435–443 (2015).
 92. Ruviaro, C. F. et al. Economic and environmental feasibility of beef production in different feed management systems in the Pampa biome, southern Brazil. *Ecol. Indic.* **60**, 930–939 (2016).
 93. Dick, M., Da Silva, M. A. & Dewes, H. Mitigation of environmental impacts of beef cattle production in southern Brazil - evaluation using farm-based life cycle assessment. *J. Clean. Prod.* **87**, 58–67 (2015).
 94. Lesnoff, M. *DynMod: a Tool for Demographic Projections of Tropical Livestock Populations Under Microsoft Excel, User's Manual - Version 1* (CIRAD, Montpellier, Cedex; ILRI, Nairobi, Kenya, 2008).
 95. Broom, D. M., Galindo, F. A. & Murgueitio, E. Sustainable, efficient livestock production with high biodiversity and good welfare for animals. *Proc. R. Soc. B* **280**, 20132025 (2013).
 96. Junior, C. C. et al. Brazilian beef cattle feedlot manure management: a country survey. *J. Anim. Sci.* **91**, 1811–1818 (2013).
 97. Garnsworthy, P. C. The environmental impact of fertility in dairy cows: a modelling approach to predict methane and ammonia emissions. *Anim. Feed Sci. Technol.* **112**, 211–223 (2004).
 98. Collins, A. L. & Zhang, Y. Exceedance of modern 'background' fine-grained sediment delivery to rivers due to current agricultural land use and uptake of water pollution mitigation options across England and Wales. *Environ. Sci. Policy* **61**, 61–73 (2016).
 99. Chadwick, D. et al. Manure management: implications for greenhouse gas emissions. *Anim. Feed Sci. Technol.* **166–167**, 514–531 (2011).
 100. *Organic Dairy Cows: Milk Yield and Lactation Characteristics in Thirteen Established Herds and Development of a Herd Simulation Model for Organic Milk Production* Project Report OF0170 (DEFRA, 2000); <https://randd.defra.gov.uk/Default.aspx?Menu=Menu&Module=More&Location=None&Completed=0&ProjectID=8431>
 101. Wilkinson, J. M. Re-defining efficiency of feed use by livestock. *Animal* **5**, 1014–1022 (2011).
 102. Webb, J., Audsley, E., Williams, A., Pearn, K. & Chatterton, J. Can UK livestock production be configured to maintain production while meeting targets to reduce emissions of greenhouse gases and ammonia? *J. Clean. Prod.* **83**, 204–211 (2014).
 103. de Ponti, T., Rijk, B. & van Ittersum, M. K. The crop yield gap between organic and conventional agriculture. *Agric. Syst.* **108**, 1–9 (2012).
 104. Gerber, P., Vellinga, T., Opio, C., Henderson, B. & Steinfeld, H. *Greenhouse Gas Emissions from the Dairy Sector: A Life Cycle Assessment* (Food and Agriculture Organization of the United Nations: 2010); <http://www.fao.org/docrep/012/k7930e/k7930e00.pdf>
 105. Brown, K. et al. *UK Greenhouse Gas Inventory, 1990 to 2010: Annual Report for Submission under the Framework Convention on Climate Change* (DEFRA, 2012); https://uk-air.defra.gov.uk/assets/documents/reports/cat07/1204251149_ukghgi-90-10_main_chapters_issue2_print_v1.pdf
 106. Misselbrook, T. H., Sutton, M. A. & Scholefield, D. A simple process-based model for estimating ammonia emissions from agricultural land after fertilizer applications. *Soil Use Manag.* **20**, 365–372 (2006).
 107. Misselbrook, T. H., Gilhespy, S. L., Cardenas, L. M., Williams, J. & Dragosits, U. *Inventory of Ammonia Emissions from UK Agriculture 2015: DEFRA Contract Report (SCF0102)* (DEFRA, 2016); https://uk-air.defra.gov.uk/library/reports?report_id=928
 108. Vellinga, T. V. et al. *Methodology Used in FeedPrint: a Tool Quantifying Greenhouse Gas Emissions of Feed Production and Utilization* Report 674 (Wageningen UR Livestock Research, 2013).
 109. Anthony, S., Quinn, P. & Lord, E. Catchment scale modelling of nitrate leaching. *Asp. Appl. Biol.* **46**, 23–32 (1996).
 110. Wang, L. et al. The changing trend in nitrate concentrations in major aquifers due to historical nitrate loading from agricultural land across England and Wales from 1925 to 2150. *Sci. Total Environ.* **542**, 694–705 (2016).
 111. Davison, P. S., Lord, E. I., Betson, M. J. & Strömqvist, J. PSYCHIC – A process-based model of phosphorus and sediment mobilisation and delivery within agricultural catchments. Part 1: Model description and parameterisation. *J. Hydrol.* **350**, 290–302 (2008).
 112. Koponen, K. & Soimakallio, S. Foregone carbon sequestration due to land occupation - the case of agro-bioenergy in Finland. *Int. J. Life Cycle Assess.* **20**, 1544–1556 (2015).
 113. Guo, L. B. & Gifford, R. M. Soil carbon stocks and land use change: a meta analysis. *Glob. Change Biol.* **8**, 345–360 (2002).

Acknowledgements

We are grateful for funding from the Cambridge Conservation Initiative Collaborative Fund and Arcadia, the Grantham Foundation for the Protection of the Environment, the Kenneth Miller Trust, the UK-China Virtual Joint Centre for Agricultural Nitrogen (CINAG, BB/N013468/1, financed by the Newton Fund via BBSRC and NERC), BBSRC (BBS/E/C/00010330), DEVIL (NE/M021327/1), U-GRASS (NE/M016900/1), Soils-R-GREAT (NE/P019455/1), N-Circle (BB/N013484/1), BBSRC Soil to Nutrition (S2N) strategic programme (BBS/E/C/00010330), UNAM-PAPIIT (IV200715), the Belmont Forum/FACEE-JPI (NE/M021327/1 'DEVIL') and the Cambridge Earth System Science NERC DTP (NE/L002507/1); A.B. is supported by a Royal Society Wolfson Research Merit award. We thank F. Brendrup, E. Caton, A. Dobermann, T. J. Florindo, E. Fonte, O. Leyser, A. Mazzetto, J. Murthwaite, F. P. Kamali, R. Olea-Perez, S. Ramsden, C. Ruviaro, J. Storkey, B. Strassburg, M. Toplioff, J. N. V. da Silva, D. Williams, X. Yan and Y. Zhang for advice, data or analysis, and K. Willott for much practical support.

Author contributions

A.B., T.A., H.B., D.C., D.E., R.F., P.G., R.G., P.S., H.W., A.W. and R.E. designed the study and performed the research; D.M.B., A.C., J.C., T.F., E.G., A.G.-H., J.H.-M., M.H., F.H., A.L., T.M., B.P., B.I.S., T.T., J.V. and E.z.E. contributed and analysed data and results; and all authors contributed substantially to the analysis and interpretation of results and writing of the manuscript.

Competing interests

The authors declare no competing interests.

Additional information

Supplementary information is available for this paper at <https://doi.org/10.1038/s41893-018-0138-5>.

Reprints and permissions information is available at www.nature.com/reprints.

Correspondence and requests for materials should be addressed to A.B.

Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.